

## BAGene product line

### Literatur / References / Bibliographie / Bibliografia

Stand / Issued / Edition / Pubblicata il: 2014-07

#### ABO-TYPE

**Ferguson-Smith MA**, Aitken DA, Turleau C, de Grouchy J: Localisation of the human ABO: Np-1: AK-1 linkage group by regional assignment of AK-1 to 9q34. Hum Genet 1976;34:35-43.

**Yamamoto F**, Clausen H, White T, Marken J, Hakamori S: Molecular genetic basis of the histo-blood group ABO system. Nature 1990; 345:229-33.

**Yamamoto F**, Hakamori S: Sugar-nucleotide donor specificity of histo-blood group A and B transferases is based on amino acid substitutions. J Biol Chem 1990;265: 19257-62.

**Yamamoto F**, McNeill PD, Yamamoto M, Hakomori S, Harris T, Judd WJ, Davenport RD: Molecular genetic analysis of the ABO blood group system: 1. Weak subgroups: A<sup>3</sup> and B<sup>3</sup> alleles. Vox Sang 1993;64:116-9.

**Yamamoto F**, McNeill PD, Yamamoto M, Hakomori S, Harris T: Molecular genetic analysis of the ABO blood group system: 3. A<sup>x</sup> and B<sup>(A)</sup> alleles. Vox Sang 1993;64:171-4.

**Chester MA**, Olsson ML. The ABO blood group gene: A locus of considerable genetic diversity. Transfus Med Rev 2001; 15:177-200.

**Yip SP**. Sequence variation at the human ABO locus. Ann Hum Genet 2002;66:1-27.

**Gassner C**, SchmarDA A, Nussbaumer W, Schönitzer D: ABO Glycosyltransferase Genotyping by polymerase chain reaction using sequence-specific primers. Blood 1996; 88:1852-6.

**Müller TH**, Hallensleben M, Schunter F, Blasczyk R: Molekulargenetische Blutgruppendiagnostik. Dt Ärztebl 2001;98:A 317-322 [Heft 6].

**Seltsam A**, Hallensleben A, Kollmann A, Burkhart J, Blasczyk R. Systematic analysis of the ABO gene diversity within exons 6 and 7 by PCR-screening revealed new ABO alleles. Transfusion 2003; 43:428-39.

**Olsson ML**, Irshaid NM, Hosseini-Maaf B, et al. Genomic analysis of clinical samples with serologic ABO blood grouping discrepancies: Identification of 15 novel A and B subgroup alleles. Blood 2001; 98:1585-93.

**Seltsam A**, Hallensleben M, Kollmann A, Blasczyk R: The nature of diversity and diversification at the ABO locus. Blood 2003; 102:3035-42.

**Seltsam A**, Das Gupta C, Wagner FF, Blasczyk R. Non-deletional ABO\*O alleles express weak blood group A phenotypes. Transfusion 2005; 45:359-65.

#### RH-TYPE, Partial D-TYPE, Weak D-TYPE, D Zygoty-TYPE

**Chérif-Zahar B**, Localization of the human Rh blood group gene structure to chromosome region 1p34.3 – 1p36.1 by in situ hybridisation. Hum Genet 1991; 86: 398-400.

**Blunt T**, Daniels G, Carritt B: Serotype switching in a partially deleted *RHD* gene. Vox Sang 1994; 67:397-401.

**Rouillac C**, Colin Y, Hughes-Jones NC, et al.: Transcript analysis of D category phenotypes predicts hybrid Rh D-CE-D proteins associated with alteration of D epitopes. *Blood* 1995; 85:2937-44.

**Rouillac C**, Le Van Kim C, Beolet M, Cartron J-P, Colin Y: Leu110Pro substitution in the RhD polypeptide is responsible for the DVII category blood group phenotype. *Am J Hematol* 1995; 49:87-8.

**Mouro I**, Colin Y, Sistonen P, Le Pennec PY, Cartron J-P, Le Van Kim C: Molecular basis of the RhC<sup>w</sup> (Rh8) and RhC<sup>x</sup> (Rh9) blood group specificities. *Blood* 1995; 86:1196-201.

**Beckers EAM**, Faas BHW, Simsek S, et al.: The genetic basis of a new partial D antigen: D<sup>DBT</sup>. *Br J Haematol* 1996; 93:720-7.

**Gassner C**, et al.: RHD/CE typing by polymerase chain reaction using sequence-specific primers *Transfusion* 1997; 37:1020-6.

**Okada H**, Kawano M, Iwamoto S, Tanaka M, Seno T, Okubo Y, Kajii E: The *RHD* gene is highly detectable in RhD-negative Japanese donors. *J Clin Invest* 1997; 100:373-379

**Wagner FF**, Gassner C, Müller TH, Schönitzer D, Schunter F, Flegel WA: Three molecular structures cause Rhesus D category VI phenotypes with distinct immunohematologic features. *Blood* 1998; 91:2157-68.

**Legler TJ**, Maas JH, Blaschke M, Malekan H, Ohto R, Lynen R, Bustami N, Schwartz DWM, Mayr WR, Köhler M, Panzer S: *RHD* genotyping in weak D phenotypes by multiple polymerase chain reactions. *Transfusion* 1998; 38:434-40.

**Omi T**, Takahashi J, Tsudo N, et al.: The genomic organization of the partial D category DVa : the presence of a new partial D associated with the DVa phenotype. *Biochem Biophys Res Commun* 1999; 254:786-94.

**Wagner FF**, Gassner C, Müller TH, Schönitzer D, Schunter F, Flegel WA: Molecular basis of weak D phenotypes. *Blood* 1999; 93:385-93.

**Flegel WA**, Wagner FF: Molecular genetics of *RH*. *Vox Sang* 2000;78(suppl 2): 109-15.

**Wagner FF**, Flegel WA: *RHD* gene deletion occurred in the *Rhesus box*. *Blood* 2000; 95:3662-8.

**Avent ND**, Reid ME: The Rh blood group system: a review. *Blood* 2000; 95:375-87.

**Legler TJ**, Wiemann V, Ohto H, Matuda I, Obara T, Uchikawa M, Köhler M: D<sup>Va</sup> Category Phenotype and Genotype in Japanese Families. *Vox Sang* 2000; 78:194-7.

**Wagner T**, Resch B, Legler TJ, Mossier C, Helmberg W, Köhler M, Lanzer G: Severe HDN due to anti-Ce that required exchange transfusion. *Transfusion* 2000; 40:571-4.

**Wagner FF**, Frohmajer A, Ladewig B, Eicher Ni, Lonicer CB, Muller TH, Siegel MH, Flegel WA: Weak D alleles express distinct phenotypes. *Blood* 2000; 95:2699-708.

**Singleton BK**, Green CA, Avent ND, Martin PG, Smart E, Daka A, Narter-Olaga EG, Hawthorne LM, Daniels G: The presence of an *RHD* pseudogene containing a 37 base pair duplication and a nonsense mutation in Africans with the RhD-negative blood group phenotype. *Blood* 2000; 95:12-18.

**Wagner FF**, Fromajer A, Flegel WA: *RHD* positive haplotypes in D negative Europeans. *BMC Genetics* 2001; 2:10.

**Müller TH**, Hallensleben M, Schunter F, Blasczyk R: Molekulargenetische Blutgruppendiagnostik. *Dt Arztebl* 2001; 98:A 317-322 [Heft 6].

**Shao CP**, Maas JH, Su YQ, Köhler M, Legler TJ: Molecular background of RhD-positive, D-negative, Del and weak D phenotypes in Chinese. *Vox Sang* 2002; 83:156-61.

**Wagner FF**, Eicher NI, Jørgensen JR, Lonicer CB, Flegel WA: DNB: a partial D with anti-D frequent in Central Europe. *Blood* 2002; 100:2253-6.

**Wagner FF**, Ladewig B, Angert KS, Heymann GA, Eicher NI, Flegel WA: The *DAU* allele cluster of the *RHD* gene. *Blood* 2002; 100:306-11.

**Perco P**, Shao CP, Mayr WR, Panzer S, Legler TJ: Testing for the D zygosity with three different methods revealed altered Rhesus boxes and a new weak D type. *Transfusion* 2003; 43:335-9.

## **KKD-TYPE**

**Lee S**, Naime DS, Reid ME, Redman CM: Molecular basis for the high-incidence antigens of the Kell blood group system. *Transfusion* 1997; 37:1117-22.

**Lee S**, Wu X, Reid M, Zelinski T, Redman C: Molecular basis of the Kell (K1) phenotype. *Blood* 1995; 85:912-6.

**Olives B**, Neau P, Bailly P, Hediger MA, Rousselet G, Cartron JP, Ripoche P: Cloning and functional expression of a urea transporter from human bone marrow cells. *J Biol Chem* 1994; 269:31649.

**Lucien N**, Sidoux-Walter F, Olives B, Moulds J, Le Pennec PY, Cartron JP, Bailly P: Characterization of the gene encoding the human Kidd blood group/urea transporter protein. Evidence for splice site mutations in Jknull individuals. *J Biol Chem* 1998; 273:12973-80.

**Cartron JP**, Ripoche P: Urea transport and Kidd blood groups. *Transfus Clin Biol* 1995; 2:309.

**Olives B**, Merriman M, Bailly P, Bain S, Barnett A, Todd J, Cartron JP, Merriman T: The molecular basis of the Kidd blood group polymorphism and its lack of association with type 1 diabetes susceptibility. *Hum Mol Genet* 1997; 6:1017.

**Neote K**, Mak JY, Kolakowski LFJ, Schall TJ: Functional and biochemical analysis of the cloned Duffy antigen: identity with the red blood cell chemokine receptor. *Blood* 1994; 84:44-52.

**Chitnis CE**, Chaudhuri A, Horuk R, Pogo AO, Miller LH: The domain on the Duffy blood group antigen for binding *Plasmodium vivax* and *P. knowlesi* malarial parasites to erythrocytes. *J Exp Med* 1996; 184:1531-6.

**Chaudhuri A**, Zbrzezna V, Polyakova J, Pogo AO, Hesselgesser J, Horuk R: Expression of the Duffy antigen in K562 cells. Evidence that it is the human erythrocyte chemokine receptor. *J Biol Chem* 1994; 269:7835.

**Chaudhuri A**, Polyakova J, Zbrzezna V, Williams K, Gulati S, Pogo AO: Cloning of glycoprotein D cDNA, which encodes the major subunit of the Duffy blood group system and the receptor for the *Plasmodium vivax* malaria parasite. *Proc Natl Acad Sci USA* 1993; 90:10793.

**Tournamille C**, Le van Kim C, Gane P, Cartron JP, Colin Y: Molecular basis and PCR-DNA typing of the Fya/Fyb blood group polymorphism. *Hum Genet* 1995; 95:407-10.

**Iwamoto S**, Omi T, Kajii E, Ikemoto S: Genomic organization of the glycoprotein D gene: Duffy blood group Fya/Fyb alloantigen system is associated with a polymorphism at the 44- amino acid residue. *Blood* 1995;85:622.

**Chaudhuri A**, Polyakova J, Zbrzezna V, Pogo AO: The coding sequence of Duffy blood group gene in humans and simians: restriction fragment length polymorphism, antibody and malarial parasite specificities, and expression in nonerythroid tissues in Duffy- negative individuals. *Blood* 1995; 85:615.

**Mallinson G**, Soo KS, Schall TJ, Pisacka M, Anstee DJ: Mutations in the erythrocyte chemokine receptor (Duffy) gene: the molecular basis of the Fya/Fyb antigens and identification of a deletion in the Duffy gene of an apparently healthy individual with the Fy(a-b-) phenotype. *Br J Haematol* 1995; 90:823-9.

**Tournamille C**, Colin Y, Cartron JP, Le van Kim C: Disruption of a GATA motif in the Duffy gene promoter abolishes erythroid gene expression in Duffy-negative individuals. *Nat Genet* 1995; 10:224-8.

**Tournamille C**, Le van Kim C, Gane P, Le Pennec PY, Roubinet F, Babinet J, Cartron JP, Colin Y: Arg89Cys substitution results in very low membrane expression of the Duffy antigen/receptor for chemokines in Fy(x) individuals. *Blood* 1998; 92:2147-56.

**Miller SA**, Dykes DD, Polesky HF: A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res* 1988; 16:1215.

**Rožman P**, Dovč T, Gassner C: Differentiation of autologous ABO, RHD, RHCE, KEL, JK, and FY blood group genotypes by analysis of peripheral blood samples of patients who have recently received multiple transfusions. *Transfusion* 2000; 40:936-42.

## **MNS-TYPE**

**Landsteiner K**, Levine P: A new agglutinable factor differentiating individual human bloods. *Proc Soc exp Biol* 1927; 24:600.

**Walsh RJ**, Montgomery C: A new human isoagglutinin subdividing the MN blood groups. *Nature* 1947; 160:504.

**Shih MC**, Yang LH, Wang NM, Chang JG: genomic typing of human red cell Miltenberger glycoporphins in a Taiwanese population. *Transfusion* 2000; 40:54-61.

**Storry JR**, Reid ME, Fetis S, Huang CH: Mutations in *GYPB* exon 5 drive the S-s-U<sup>var</sup> phenotype in persons of African descent: implications for transfusion. *Transfusion* 2003; 43:1738-47.

## **HPA-TYPE**

**Ballem PJ**, Buskard NA, Decary F, et al.: Post-transfusion purpura secondary to passive transfer of anti-PI<sup>A1</sup> by blood transfusion. *Br J Hematol* 1987; 66:113-4.

**Mueller-Eckhardt C**, Kiefel V, Grubert A, et al.: 348 cases of suspected neonatal alloimmune thrombocytopenia. *Lancet* 1989; 1:363-6.

**Panzer S**, Kiefel V, Bartram CR, et al.: Immune thrombocytopenia more than a year after allogeneic marrow transplantation due to antibodies against donor platelets with anti-PI<sup>A1</sup> specificity: evidence for a host derived immune reaction. *Br J Hematol* 1989; 71:259-64.

**Mueller-Eckhardt C**, Kiefel V, Santoso S: Review and update of platelet alloantigen systems. *Transfus Med Rev* 1990; 4:98-109.

**Santoso S**, Kiefel V: Human platelet specific alloantigens: update. *Vox Sang* 1998; 74 (Suppl):249-53.

**Jau-Yi L**, Ying-Ju C, Hui-Yu H, Jeong-Shi L, Cheng-Hwai T: PCR with sequence-specific primer-based simultaneous genotyping of human platelet antigen-1 to -13w. *Transfusion* 2002; 42:1089-95.

## **HNA-TYPE**

**Carl B**, Kroll H, Bux J, Bein G, Santoso S: B-lymphoblastoid cell lines as a source of reference DNA for human platelet and neutrophil antigen genotyping. *Transfusion* 2000; 40:62-68.

**Steffensen R**, Gülen T, Varming K, Jersild C: FcγRIIIB polymorphism: evidence that NA1/NA2 and SH are located in two closely linked loci and that the SH allele is linked to the NA1 allele in the Danish population. *Transfusion* 1999; 39:593-598.

**Greinacher A**, Wesche J, Hammer E, Füll B, Völker U, Reil A, Bux J: Characterization of the human neutrophil alloantigen-3a. *Nature Medicine* 16, 45-48 (2009)

**Sachs UJH**, Chavakis T, Fung L, Lohrenz A, Bux J, Reil A, Ruf A, Santoso S: Human alloantibody anti-Mart interferes with Mac-1-dependent leukocyte adhesion. *Blood* 2004; 104,3:727-34.

**Sachs UJH**, Reil A, Bauer C, Bux J, Bein G, Santoso S: Genotyping of human neutrophil antigen-5a (Ond). *Transfus Med* 2005; 15:115-117.

## **General**

**Green and Sambrook**, 2012. *Molecular Cloning: A Laboratory Manual*. New York: Cold Spring Harbour Laboratory

**Beutler E** et al., 1990. *BioTechniques* 9:166

**Olerup O**, Zetterquist H. HLA-DR typing by PCR amplification with sequence-specific primers (PCR-SSP) in 2 hours: an alternative to serological DR typing in clinical practice including donor-recipient matching in cadaveric transplantation. *Tissue Antigens* 1992; 39:225-35.

**Olerup O**, Zetterquist H. DR "low-resolution" PCR-SSP typing - a correlation and an update. *Tissue Antigens* 1993; 1:55-6.

**Chen EY**, Liao YC, Smith DH, Barrera-Saldana HA, Gelinis RE, Seeburg PH. The human growth hormone locus: nucleotide sequence, biology, and evolution. *Genomics* 1989; 4:479.



**BAG Health Care GmbH**

Amtsgerichtsstraße 1-5  
35423 Lich/Germany

Tel.: +49 (0) 6404 / 925-0

Fax: +49 (0) 6404 / 925-250

[www.bag-healthcare.com](http://www.bag-healthcare.com)

[info@bag-healthcare.com](mailto:info@bag-healthcare.com)

**Auftragsannahme/Ordering:**

Tel.: +49 (0) 6404 / 925-450

Fax: +49 (0) 6404 / 925-460

[verkauf@bag-healthcare.com](mailto:verkauf@bag-healthcare.com)

**Customer Service:**

Tel.: +49 (0) 6404 / 925-125

Fax: +49 (0) 6404 / 925-421

[service@bag-healthcare.com](mailto:service@bag-healthcare.com)